




Official Journal of the European Paediatric Neurology Society



Original article

Pyridoxine dependent epilepsy: Is late onset a predictor for favorable outcome? ☆



R.L.P. de Rooy ^a, F.J. Halbertsma ^b, E.A. Struijs ^c, F.J. van Spronsen ^d,
 R.J. Luning ^e, H.M. Schippers ^f, P.M. van Hasselt ^g, B. Plecko ^h,
 G. Wohlrab ^h, S. Whalen ⁱ, J.F. Benoist ^j, S. Valence ^k, P.B. Mills ^l,
 L.A. Bok ^{b,*}

^a Department of Pediatrics, Zuyderland Hospital, Heerlen, The Netherlands

^b Department of Pediatrics, Maxisma Medical Center, Veldhoven, The Netherlands

^c Metabolic Unit, Department of Clinical Chemistry, VU University Medical Center, Amsterdam, The Netherlands

^d Department of Metabolic Diseases, Beatrix Children's Hospital, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

^e Department of Child Neurology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

^f Department of Neurology, Sint Antonius Ziekenhuis, Nieuwegein, Utrecht, The Netherlands

^g Department of Pediatric Metabolic Diseases, Wilhelmina Children's Hospital, University Medical Center (UMC) Utrecht, Utrecht, The Netherlands

^h Division of Neurology, Children's Hospital, University of Zurich, Zurich, Switzerland

ⁱ UF de génétique clinique, APHP, Hôpital Armand Trousseau, Paris, France

^j Centre de Référence des Maladies Héritaires du Métabolisme, Service de Biochimie-Hormonologie, Hôpital Robert Debré, Paris, France

^k Department of Child Neurology, APHP, Armand Trousseau Hospital, Paris, France

^l Clinical and Molecular Genetics Unit, UCL Institute of Child Health, London, United Kingdom

ARTICLE INFO

Article history:

Received 21 March 2017

Received in revised form

6 March 2018

Accepted 25 March 2018

Keywords:

Pyridoxine dependent epilepsy

Late onset

ABSTRACT

Aim: In pyridoxine dependent epilepsy (PDE), patients usually present with neonatal seizures. A small subgroup is characterized by late-onset beyond 2 months of age. We aim to analyze the observation of relatively good cognitive outcome in this subgroup of late-onset PDE patients. **Methods:** We retrospectively analyzed data from four metabolically and genetically confirmed late-onset patients with PDE due to antiquitin (ALDH7A1) deficiency. Data were analyzed regarding ALDH7A1 mutations, alpha-Amino adipic semialdehyde (α -AASA) and pipercolic acid (PA) levels, medication during pregnancy, delivery, treatment delay, amount of seizures, pyridoxine dose, adjuvant therapy and findings on brain MRI.

Results: Results showed that three patients had relatively good outcome (IQ 80–97), while one patient did not undergo formal testing and was considered mildly delayed. We were unable to find a clear association between the above-mentioned variables and cognitive outcome, although a less severe genotype may be present in three patients, and maternal medication could be accountable for better outcome in two patients.

☆ Late-onset PDE is a phenotype associated with relatively good outcome.

* Corresponding author. PO Box 7777, 5500MB, Veldhoven, The Netherlands.

E-mail address: l.bok@mmc.nl (L.A. Bok).

<https://doi.org/10.1016/j.ejpn.2018.03.009>

1090-3798/© 2018 European Paediatric Neurology Society. Published by Elsevier Ltd. All rights reserved.

Interpretation: We suggest that favorable outcome in late onset PDE might be explained by a combination of factors. A yet unknown protective factor, different genetic variations, functional variation and secondarily variation in treatment regimens and absence of neonatal seizure induced brain damage.

© 2018 European Paediatric Neurology Society. Published by Elsevier Ltd. All rights reserved.

1. Introduction

Most cases of pyridoxine dependent epilepsy (PDE) are due to mutations in the gene, located on 5q31,^{1,2} which encodes for antiquitin, the α -AASA dehydrogenase enzyme within the lysine degradation pathway. Antiquitin deficiency results in alpha-amino adipic semialdehyde (α -AASA) and pipercolic acid (PA) accumulation in blood plasma, urine and cerebrospinal fluid. Most patients present with intractable seizures in the neonatal period. Rarely PDE patients present after the neonatal period. In the Netherlands, the incidence is 1 in 276.000 new-borns.^{3,4} The incidence of late-onset PDE is an estimated 10% of PDE, so roughly 1 in 2.500.000. In 2001, Baxter reported a more favorable outcome in late onset PDE. However, PDE could, at that time, not be confirmed by metabolic and/or DNA testing and thus not differentiated from pyridoxine responsive seizures for example.⁵ Hence it is unclear whether these patients were indeed suffering from antiquitin deficiency, or from a different form of pyridoxine dependent epilepsy.

Diagnosis can be blurred by non-neuronal systemic symptoms or ambiguous pyridoxine response. Currently, measurement of alpha-amino adipic semialdehyde in urine supports the diagnosis, followed by molecular investigations of the Antiquitin-gene.^{2,3}

Treatment of PDE is primarily based on a daily pyridoxine dose of 10–30 mg/kg/day for infants, while for neonatal patients a daily maximum of 200 mg is recommended.^{5,6} Adjuvant treatment options include dietary lysine restriction and arginine supplementation.^{7,8}

Without adequate treatment, PDE is a potentially lethal condition. Despite being seizure free, 75% of the patients treated with pyridoxine are mentally low average or even retarded (IQ < 85 or 50% IQ < 70).^{7,12} Prognosis in PDE is probably influenced by several different factors and cannot be predicted easily. One of these prognostic factors might be the patient age at seizure onset, which has been reported previously.⁵

2. Methods

We retrospectively collected data from specific patients with a metabolically (increased α -AASA in urine) as well as genetically confirmed PDE with a late onset, identified in the PDE network database. The PDE network database is a non-formal database, based on intercollegial communication between metabolic PDE centers, including the Dutch centers. Late onset was defined as onset of seizures beyond 1 month of age. Data

were obtained, if applicable from patients' physicians through a questionnaire concerning outcome and possible predicting cofactors (pregnancy and labor complications, onset of seizures, metabolic and genetic testing, treatment, outcome and MRI as well as EEG findings). Outcome was measured as IQ or development by age appropriate assessment. Bayley Scale of Infant and Toddler Development -Dutch adaptation (Bayley-II-NL) or Snijders-Oomen Non-verbal intelligence test (revised) (SON-R 81), a standardized Dutch intelligence test and the age appropriate assessment by 'van Wiechenschema' (vWS), which is based on development of healthy full-term-born children.^{10–12} Data were compared with those in early onset PDE patients as reported in the literature.

3. Results

In the PDE network database, four patients were identified that met the inclusion criteria. The main outcome IQ was available in three patients (81, 95, and 97) and appeared higher than expected based on average IQ in early onset PDE patients (IQ $72 \pm$ SD 19)¹².

One patient did not undergo validated developmental testing, but was considered having mild-moderate delay from a clinical perspective, as well as from attending normal school with extra support.¹² Major behavioral disturbances or autistic spectrum disorders were not observed in any patient. All patients were seizure free.

Results are shown in Table 1.

MRI was normal in three of our four patients. However, patient 1 showed a hypoplastic isthmus of the corpus callosum and moderately widened cerebrospinal fluid spaces.

4. Discussion

The main observation is that our study confirms that cognitive development may be better in late onset PDE, as three out of four PDE patients had IQ levels within the normal range while in typical early onset PDE, 75% of children are mentally delayed.¹² Follow-up tests to further evaluate cognitive development over time in our patients are not yet available. The role of various factors which might explain good outcome in late onset PDE are discussed below.

A genotype phenotype relation was previously assumed, but has not been established in PDE.^{2,13} Interestingly, three patients presented here had uncommon mutations with unknown residual enzyme activity. Although speculative, residual enzyme activity might be preserved, which could

Table 1 – Patient specific clinical characteristics.

	Patient 1	Patient 2	Patient 3	Patient 4
Sex	Male	Female	Female	Male
Medication during pregnancy	None	Mercaptopurine 75 mg	None	None
Abnormal fetal movements	None	Hiccups	None	None
Vitamins during pregnancy	Vitamin B12 IM 1000 µg monthly	Vitamin B6 2 mg (=ADD) ^a	Folic Acid	None
Pregnancy complications	Multiple pregnancy (Twins)	Maternal Crohn's Disease	Placental problems	Glucocorticoid for preterm labour at 26 weeks of gestation
Labor complications	None	Meconium staining	None	Preterm, 31 gestational weeks
Apgar	9/10	7/9	–/–	8/10
Neonatal period	Normal	Normal	Normal	Transient neonatal respiratory distress Transient enteropathy
First seizures	10th week	20th week	9th week	3 months
First pyridoxine	13th week	14th month	12th week	3 months
α -AASA in urine at diagnosis (mmol/mol creatinin)	101	9.9	13	Not available
Mutation ^a	c.1279G > C c.1279G > C	c.346C > T c.1432T > A	c.512C > T c.512C > T	c.1364T > C c.1364T > C
Amino acid change ^a	p.Glu427Gln p.Glu427Gln	p.Gln116Stop p.Cys478Ser	p.Ala199Val p.Ala199Val	p.Leu455Pro p.Leu455Pro
Pipecolic acid in plasma (µmol/l)	10.9	3.0	4.91	16.4
Outcome: current age	4years	4 years	10 years	4 years
Pyridoxine dose	100 mg	300 mg (200 morning/ 100 evening)	300 mg	250 mg
Current antiepileptic drugs	None	None	None	n.a. ^a
Lysine restricted diet	Yes, starting at 2.5 years	No	No	No
IQ/MD	Bailey 97 (age 2)	Bailey 95 (age 2)	SON-R 81; concentration deficits	Attends normal school with extra support
Behavior	Normal	Normal	Normal	Shy
MRI: at diagnosis	Hypoplastic isthmus Moderately widened CSF ^a spaces	Normal	Normal	Normal
EEG at diagnosis	Spike waves centroparietal	Normal background, no epileptic discharges	Seizure pattern O1/O2 ^a , independent. Normal background activity	Spikewaves right temporo-parietal
EEG at follow-up	Normal	Normal	Normal	Normal

ADD = advised daily dose; CSF = cerebrospinal fluid. n.a. = not available. O = occipital electrode. IQ = intelligence quotient. MD = mental development. SON-R=Snijders-Oomen Non-verbal intelligence test (revised) Reference range for urinary α -AASA: <1 mmol/mol creatinin Reference range for plasma Pipecolic Acid: 3.75–10.8 µmol/L (in first week of life); 0.7–2.46 µmol/L (after first week).

^a The reference sequence used for reporting DNA variants is NM_001182.4.

explain late onset and better prognosis. Patient 1, however, was homozygous for the common Dutch mutation c.1279G > C (formerly designated as c.1195G > C) reported as having an unfavorable outcome.¹²

Comparing PA levels at time of diagnosis of early onset patients to the ones observed here, two patients (2 and 3) had low to medium PA values.³ Patients 1 and 4 showed a relatively high value of plasma PA, which could be a reflection of higher lysine intake or low residual enzyme activity. However, these patient's outcome in terms of intelligence were not worse compared to others in this study.

In early onset PDE, α -AASA values range from 4.0 to 29 mmol/mol creatinine in urine at the time of diagnosis.³ Again, in two patients, the α -AASA levels in urine were moderately elevated. One patient (patient 1) showed a strongly increased urinary α -AASA level of 101 mmol/mol creatinine at diagnosis, but nevertheless the patient had a normal cognitive outcome. In patient 4, urinary α -AASA concentration was not available.

Delay in treatment was regarded as the time between seizure onset and start of pyridoxine treatment and ranged from none to nine months. Delay in treatment in this cohort showed notable differences (Table 1) that could not be related to the favorable outcome in our patients. This is in disagreement with the findings of a UK national survey that showed treatment delays varying from 1 day to 53 weeks in early onset PDE, associating longer delay with disadvantageous prognosis.⁵

The received pyridoxine dose of patients in this study, as shown in Table 1, was varying from 6 to 22 mg/kg/day, based on an age-dependent estimated weight, either in a single daily dose or in two uneven daily doses (e.g. 200 mg in the morning and 100 mg in the evening). Currently, the recommended dose of pyridoxine is 15–30 mg/kg/day in early infancy, reducing to doses ranging from 0.5 to 2.0 mg/kg/day in later life.⁵ This recommended dose was based on early-onset patients however. The patients we studied were treated with pyridoxine dosages within this recommended range. Higher doses might

be associated with better behavioral and electroencephalographical outcome.⁵ This however does not count for the patients included in this study, as they received relatively low doses.

Three of the patients in our study received monotherapy with pyridoxine, whereas adjuvant treatment with lysine restriction and arginine supplementation have been suggested.^{6,7} Patient 1 started lysine restriction at the age of 2.5 years, whilst IQ-testing was performed at 2 years of age. Based on these data we cannot conclude that there might be a positive effect of lysine restriction on IQ.

There were differences with respect to medication during pregnancy. The mother of patient 2 used mercaptopurin and vitamin B6, as a therapy for Crohn's disease. Vitamin B6 was used in the advised daily dose of 2 mg. Although mercaptopurin occasionally has been associated with congenital defects in the newborn and pregnancy complications,¹⁴ the current opinion in the Netherlands is that a low dose of mercaptopurin can safely be used during pregnancy.¹⁵ Possible benefits from mercaptopurin use during pregnancy could not be found in literature. A neuroprotective effect seems very unlikely. Thus, for our patient, we could not find a clear answer on possible disadvantages or benefits resulting from maternal mercaptopurin use during pregnancy, concluding that it is highly unlikely to be of any influence on our patients' good outcome.

The use of folic acid and vitamin B12 in the mother of patient 1 and 3 respectively is noteworthy. No studies investigating the effects of maternal folic acid nor vitamin B12 use during pregnancy on epilepsy syndromes in infants were found, nor a relation with these epilepsy syndromes. Positive effects of folic acid in PDE however, have been described before.¹⁶ Positive effects of vitamin B12 use in PDE have not yet been described. Vitamin B12 and folic acid metabolism are linked in the remethylation cycle of homocysteine, but its role in PDE is not determined.

The mother of patient 4 received glucocorticoids because of possible preterm birth at 26 weeks. A relation between maternal glucocorticoid use and seizures in their child was not found in literature, and therefore cannot be held responsible for favorable outcome in late onset PDE.

Seizure control has recently been suggested as a factor in neurocognitive development. It is assumed that neonatal seizures regardless of their etiology are a great threat for neurocognitive development, especially if they occur in the first weeks of life.¹⁷ This might be applicable in this study of late-onset PDE patients. Therefore, patients with late onset of seizures and good outcome are of significant importance in unraveling the pathophysiology of PDE, which also might enhance the prognosis of all PDE patients.

The observations in this study leave room for speculations. One might speculate that there is a yet to be discovered protective biochemical –detoxification- process responsible for good outcome in late-onset PDE patients.

4.1. Study limitations

In PDE, and many other inborn-errors of metabolism, low number of patients impedes research. This certainly applies for the subgroup of late-onset PDE patients. Nevertheless, only

by sharing observations, understanding of PDE can progress. Follow-up data concerning developmental outcome were incomplete for patient 4. For this patient, only clinical observations were available, as no validated developmental test was performed. This patient is considered to be mildly delayed, given the fact he attends a normal school with extra support. From this perspective, we still think it is of added value to compare this definite late onset patient to others in this study, especially since late-onset PDE is such a rare disease.

5. Conclusion

In late onset PDE, IQ tends to be higher than in the majority of classic early onset PDE. There is a possible role in relation to outcome for genetics, as a possible favorable genotype–phenotype correlation, metabolic factors as lower AASA and lower pipercolic acid levels, adjuvant therapy as lysine restriction or maternal medication during pregnancy as folic acid. In comparison with reported cases no uniform explanation for mild outcome in these studied late-onset PDE patients could be found; except the coincidence of (prolonged) absence of neonatal seizures. There might be a yet to be discovered protective factor in the patients.

Conflict of interest

None.

REFERENCES

1. Bok LA, Struys E, Jakobs C, Been JV, Willemsen MAA. Pyridoxine Afhankelijke Epilepsie. *Ned Tijdschr Neurol en Neuroch* 2007;108:51–6.
2. Mills PB, Struys E, Jakobs C, Plecko B, Baxter P, Baumgartner M, et al. Mutations in antiquitin in individuals with pyridoxine-dependent seizures. *Nat Med* 2006;12:307–9.
3. Bok L, Struys E, Willemsen M, Been J. Pyridoxine-dependent seizures in Dutch patients: diagnosis by elevated urinary alpha-aminoadipic semialdehyde levels. *Arch Dis Child* 2007;92:687–9.
4. van Karnebeek CD, Tiebout SA, Niermeijer J, Poll-The BT, Ghani A, Coughlin 2nd CR. Pyridoxine-dependent epilepsy: an Expanding clinical spectrum. *Pediatr Neurol* 2016 Jun;59:6–12.
5. Baxter P. *Pyridoxine dependent/responsive seizures; in vitamin responsive conditions in paediatric neurology*. London: Mac Keith Press; 2001. p. 109–65.
6. Stockler S, Plecko B, Gospe Jr SM, et al. Pyridoxine dependent epilepsy and antiquitin deficiency clinical and molecular characteristics and recommendations for diagnosis, treatment and follow-up. *Mol Genet Metabol* 2011;104:48–60.
7. Van Karnebeek CD, Hartmann H, Jaggamantri S, Bok LA, Cheng B, Connolly M, Coughlin 2nd CR, Das AM, Gospe Jr SM, Jakobs C, van der Lee JH, Mercimek-Mahmutoglu S, Meyer U, Struys E, Sinclair G, Van Hove J, Collet JP, Plecko BR, Stockler S. Lysine restricted diet for pyridoxine-dependent epilepsy: first evidence and future trials. *Mol Genet Metabol* 2012 Nov;107(3):335–44.

8. Van Karnebeek CD, Jagdumantri S. Current treatment and management in pyridoxine-dependent epilepsy. *Curr Treat Options Neurol* 2015 Feb;**17**(2):335.
10. Tellegen PJ, Laros JA. SON-R 6–40. *Snijders-Oomen niet-verbale intelligentietest. I. Verantwoording*. Amsterdam: Hogrefe uitgevers; 2011.
11. Laurent de Angulo MS, Brouwers-de Jong EA, Bijlsma-Schlösser JFM, et al., editors. *Ontwikkelingsonderzoek in de jeugdgezondheidszorg: het Van Wiechenonderzoek - De Baecke-Fassaert Motoriektest. 4 e dr*. Assen: Van Gorcum; 2008.
12. Bok LA, Halbertsma FJ, Houterman S, Wevers RA, Vreeswijk C, Jakobs C, Struys E, van der Hoeven JH, Sival DA, Willemsen MA. Long-term outcome in pyridoxine-dependent epilepsy. *Dev Med Child Neurol* 2012;**54**:849–54.
13. Schärer G, Brocker C, Vasiliou V, Creadon-Swindell G, Gallagher RC, Spector E, Van Hove JL. The genotypic and phenotypic spectrum of pyridoxine-dependent epilepsy due to mutations in ALDH7A1. *J Inherit Metab Dis* 2010 Oct;**33**(5):571–81.
14. Nørgård B, Pedersen L, Christensen LA, Sørensen HT. Therapeutic drug use in women with Crohn's disease and birth outcomes: a Danish nationwide cohort study. *Am J Gastroenterol* 2007;**102**(7):1406.
15. Francella A, Dyan A, Bodian C, Rubin P, Chapman M, Present DH. The safety of 6-mercaptopurine for childbearing patients with inflammatory bowel disease: a retrospective cohort study. *Gastroenterology* 2003;**124**(1):9.
16. Gallagher RC, Van Hove JL, Schärer G, Hyland K, Plecko B, Waters PJ, Mercimek-Mahmutoglu S, Stockler-Ipsiroglu S, Salomons GS, Rosenberg EH, Struys EA, Jakobs C. Folinic acid-responsive seizures are identical to pyridoxine-dependent epilepsy. *Ann Neurol* 2009 May;**65**(5):550–6.
17. Chapman KE, Specchio N, Shinnar S, Holmes GL. Seizing control of epileptic activity can improve outcome. *Epilepsia* 2015;**56**:1482–5.